

ESSENTIAL NORMS OF TOEPLITZ OPERATORS ON BERGMAN-HARDY SPACES ON THE UNIT DISK

ARTUR MICHALAK

Abstract: For a Borel measure μ on $[0, 1]$ with $1 \in \text{supp}(\mu)$ we consider the Toeplitz operators T_f with $f \in L^\infty(\mu \otimes \lambda)$ on the space $H^2(\mu)$ consisting of all holomorphic on \mathbb{D} functions in the Lebesgue class $L^2(\mu \otimes \lambda)$ on \mathbb{D} . We show that for every Bergman-Hardy space $H^2(\mu)$ the following estimations hold: $\text{dist}(T_f, \mathcal{K}(H^2(\mu))) \geq |f(t_0)|$ if f is continuous at point $t_0 \in \mathbb{T}$, and $\text{dist}(T_f, \mathcal{K}(H^2(\mu))) = \sup\{|f(z)| : z \in \mathbb{D}\}$ if f is a bounded harmonic function on \mathbb{D} .

Keywords: Toeplitz operators, Bergman spaces

The Bergman-Hardy spaces on the disk are generalization of the weighted Bergman spaces on the disk. Informally we can say that a weighted Bergman space on the disk is generated by the measure which in polar coordinates has the form $(1 - r^2)^\alpha r dr \times \lambda$ for some $\alpha > -1$, where dr and λ are the Lebesgue measures on $[0, 1]$ and \mathbb{T} , respectively. A Bergman-Hardy space $H^p(\mu)$ on the disk is generated by the measure of the form $\mu \times \lambda$ in polar coordinates, where μ is an arbitrary positive finite Borel measure on $[0, 1]$ which do not vanish near 1. We do not have explicit formula for the Bergman kernel as well as for a reproducing kernel for all Bergman-Hardy spaces $H^2(\mu)$, and we can not apply the Berezin transform techniques (see [9], [8]) for these spaces. Our approach is more directly, we apply sequences of functions in $H^2(\mu)$ which are closely related to the Cauchy and Poisson kernels in \mathbb{D} . We show that the classical results (see [1], [9]) for Bergman and weighted Bergman spaces on the disk concerning essential norms of Toeplitz operators T_f , when f has a point of continuity on the boundary or f is harmonic, hold for every Bergman-Hardy space $H^2(\mu)$.

The paper is divided into two sections. The terminology and basic facts are explained in the first. We gather in this section basic properties of Bergman-Hardy spaces $H^p(\mu)$ for $0 < p < \infty$. In the second section we estimate the

1991 Mathematics Subject Classification: Primary 47B35

Acknowledgment. This research was mostly done when the author held a DAAD grant at Paderborn University. The research was supported in part by Komitet Badań Naukowych (State Committee for Scientific Research), Poland, grant no. 2 P03A 051 15.

essential norms of Toeplitz operators T_f if f has a point of continuity on the boundary \mathbb{T} or is harmonic.

1. Preliminaries

We start by explaining basic notation used in this paper. As usual, \mathbb{D} will stand for the open unit disk and \mathbb{T} for the unit circle in the complex plain \mathbb{C} . The Lebesgue measures on $[0, 1]$ and \mathbb{T} will be denoted by dr and λ , respectively. Throughout the paper μ will be a positive Borel measure on $[0, 1]$ with $1 \in \text{supp}(\mu)$ (the support of μ is the smallest closed set $C \subset [0, 1]$ such that $\mu(C) = \mu([0, 1])$). By $\mu \otimes \lambda$ we will denote the Borel measure on $\overline{\mathbb{D}}$ given by $\mu \otimes \lambda(A) = \mu \times \lambda(\Phi^{-1}(A))$ where $\Phi : [0, 1] \times \mathbb{T} \rightarrow \overline{\mathbb{D}}$ is given by $\Phi((r, t)) = rt$. The norm of an element $f \in L^\infty(\mu \otimes \lambda)$ will be denoted by $\|f\|_\infty$. The space of all continuous linear operators on a given Hilbert space H taking values in H equipped with the operator norm will be denoted by $\mathcal{L}(H)$ and its subspace consisting of all compact operators by $\mathcal{K}(H)$. The essential norm of an operator $T \in \mathcal{L}(H)$ is its distance to compact operators,

$$\text{dist}(T, \mathcal{K}(H)) = \inf\{\|T - S\| : S \in \mathcal{K}(H)\}.$$

For a positive finite Borel measure μ on $[0, 1]$ with $1 \in \text{supp}(\mu)$ and $0 < p < \infty$, we denote by $H^p(\mu)$ the space of all holomorphic functions $f : \mathbb{D} \rightarrow \mathbb{C}$ such that

$$\|f\| = \left(\int_{[0,1]} \int_{\mathbb{T}} |f(rt)|^p d\lambda(t) d\mu(r) + \mu(\{1\}) \sup_{0 < r < 1} \int_{\mathbb{T}} |f(rt)|^p d\lambda(t) \right)^{1/p} < \infty.$$

equipped with the norm (when $1 \leq p < \infty$) and quasi-norm (when $0 < p < 1$) defined above. There are many classical examples of spaces of holomorphic functions on \mathbb{D} that are Bergman-Hardy spaces. For example: $H^p(\delta_1)$ is the classical Hardy space $H^p(\mathbb{D})$ of holomorphic functions on \mathbb{D} (δ_1 is the Dirac measure on $[0, 1]$ concentrated at 1), the space $H^p(r dr)$ is the classical Bergman space of holomorphic functions on \mathbb{D} , the space $H^p((\alpha + 1)(1 - r^2)^\alpha r dr)$ for some $\alpha > -1$ is the classical weighted Bergman space of holomorphic functions on \mathbb{D} . The definition above seems to be artificial, but it generates spaces consisting of holomorphic functions on \mathbb{D} which are closed subspaces of $L^p(\mu \otimes \lambda)$ with a dense subset of polynomials. In order to check these facts we will need

Proposition 1. *Let μ be a positive finite Borel measure on $[0, 1]$ such that $1 \in \text{supp}(\mu)$ and $0 < p < \infty$. For every $f \in H^p(\mu)$ and $z \in \mathbb{D}$*

$$|f(z)| \leq \begin{cases} \left(\frac{4}{(1 - |z|)\mu((1 + |z|)/2, 1]} \right)^{1/p} \|f\| & \text{if } \mu(\{1\}) = 0 \\ \left(\frac{4}{(1 - |z|)\mu(\{1\})} \right)^{1/p} \|f\| & \text{if } \mu(\{1\}) > 0 \end{cases}$$

Proof. Applying the fact that $|f|^p$ is a subharmonic function we get

$$|f(z)|^p \leq \int_{\mathbb{T}} \frac{R^2 - |z|^2}{|Rt - z|^2} |f(Rt)|^p d\lambda(t) \leq \frac{R + |z|}{R - |z|} \int_{\mathbb{T}} |f(Rt)|^p d\lambda(t)$$

for every $|z| < R < 1$. Therefore, if $\mu(\{1\}) = 0$,

$$\begin{aligned} |f(z)|^p &\leq \frac{4}{(1 - |z|)\mu([(1 + |z|)/2, 1])} \int_{[(1 + |z|)/2, 1]} \int_{\mathbb{T}} |f(rt)|^p d\lambda(t) d\mu(r) \\ &\leq \frac{4 \|f\|^p}{(1 - |z|)\mu([(1 + |z|)/2, 1])}. \end{aligned}$$

If $\mu(\{1\}) > 0$,

$$|f(z)|^p \leq \frac{4 \|f\|^p}{(1 - |z|)\mu(\{1\})}.$$

■

From the proposition follows that the topology of $H^p(\mu)$ is stronger than the topology of uniform convergence on compact subsets of \mathbb{D} . Then spaces $H^p(\mu)$ are Banach for $1 \leq p < \infty$ and p -Banach for $0 < p < 1$. If $\mu(\{1\}) = 0$, then $H^p(\mu)$ is a closed subspace of $L^p(\mu \otimes \lambda)$. If $\mu(\{1\}) > 0$, then $H^p(\mu)$ coincides with $H^p(\mathbb{D})$, moreover the norms $\|\cdot\|_{H^p(\mu)}$ and $\|\cdot\|_{H^p(\mathbb{D})}$ are equivalent. For every f from the Hardy class $H^p(\mathbb{D})$

$$\sup_{0 < r < 1} \int_{\mathbb{T}} |f(rt)|^p d\lambda(t) = \int_{\mathbb{T}} |f^*(t)|^p d\lambda(t)$$

where f^* is the radial limit function of f . If $\mu(\{1\}) > 0$, we will identify $f \in H^p(\mu)$ with the function \tilde{f} on $\overline{\mathbb{D}}$ given by $\tilde{f}|_{\mathbb{D}} = f$ and $\tilde{f}|_{\mathbb{T}} = f^*$. Then the norm (p -norm) in $H^p(\mu)$ can be evaluate applying the following formula

$$\|f\| = \left(\int_{\overline{\mathbb{D}}} |f|^p d\mu \otimes \lambda \right)^{\frac{1}{p}}.$$

Hence, $H^p(\mu)$ is a closed subspace of $L^p(\mu \otimes \lambda)$ also if $\mu(\{1\}) > 0$. The space $L^p(\mu \otimes \lambda)$ is separable for every $0 < p < \infty$, it is a straightforward consequence of the Lusin theorem and the fact that the space $C(\overline{\mathbb{D}})$ of all complex continuous functions on $\overline{\mathbb{D}}$ is separable. Therefore the space $H^p(\mu)$ is separable for every μ . For every $t \in \mathbb{T}$ and $f \in H^p(\mu)$, $\|f\| = \|f_t\|$ where $f_t(z) = f(zt)$. By standard arguments concerning properties of translations (see [3], [7]) polynomials are dense in $H^p(\mu)$ for every μ . Furthermore, the closed unit ball $B_{H^p(\mu)}$ of $H^p(\mu)$ is compact in the topology of uniform convergence on compact subsets of \mathbb{D} . Consequently, the weak topology and the topology of uniform convergence on compact subsets of \mathbb{D} coincides on $B_{H^p(\mu)}$ for every $1 < p < \infty$.

2. The main results

The space $H^2(\mu)$ is a closed subspace of the Hilbert space $L^2(\mu \otimes \lambda)$. Let P be the orthogonal projection of $L^2(\mu \otimes \lambda)$ onto $H^2(\mu)$. For every $f \in L^\infty(\mu \otimes \lambda)$ the Toeplitz operator $T_f : H^2(\mu) \rightarrow H^2(\mu)$ is given by

$$T_f(h) = P(fh).$$

Since P is orthogonal, for every $g, h \in H^2(\mu)$

$$\langle g, T_f(h) \rangle = \langle P^*(g), fh \rangle = \langle g, fh \rangle.$$

In the sequel we will also need the following elementary

Fact 2. For every $\pi \geq \delta > 0$ and $R > r > 0$

$$\int_{-\delta}^{\delta} \frac{dt}{|R - re^{it}|^2} = \frac{4}{R^2 - r^2} \operatorname{arctg} \left(\frac{R+r}{R-r} \operatorname{tg} \left(\frac{\delta}{2} \right) \right).$$

Now we are ready to state our main result.

Theorem 3. Let μ be a positive finite Borel measure on $[0, 1]$ such that $1 \in \operatorname{supp}(\mu)$. If $f \in L^\infty(\mu \otimes \lambda)$ is continuous at point $t_0 \in \mathbb{T}$, then

$$\operatorname{dist}(T_f, \mathcal{K}(H^2(\mu))) \geq |f(t_0)|.$$

Proof. We can assume that $t_0 = 1$ and $f(1) > 0$. Let us take any $\varepsilon > 0$. Since f is continuous at 1, there exists an open neighborhood U of 1 such that $|f(rt) - f(1)| < \varepsilon$ for every $rt \in U$. Hence, there exists $\delta \in (0, 1)$ such that $re^{it} \in U$ for every $t \in [-\delta, \delta]$ and $r \in [1 - \delta, 1]$. Let

$$g_n(z) = \frac{z^n}{1 + \delta/n - z}.$$

Then

$$\|g_n\|^2 = \int_0^1 \int_{\mathbb{T}} \frac{r^{2n}}{|1 + \delta/n - rt|^2} d\lambda(t) d\mu(r) = \int_0^1 \frac{r^{2n}}{(1 + \delta/n)^2 - r^2} d\mu(r).$$

Since $r \leq 1$ and $(1 + \delta/n)^2 - r^2 \geq 2\delta/n$ for $0 \leq r \leq 1$,

$$\|g_n\|^2 \leq n \frac{\mu([0, 1])}{2\delta}$$

On the other hand, for every $0 < r_0 < 1$

$$\|g_n\|^2 \geq \frac{r_0^{2n}}{4} \mu([r_0, 1]).$$

It follows that $(g_n/\|g_n\|)$ is a null sequence in the topology of uniform convergence on compact subsets of \mathbb{D} . Consequently, it converges to zero in the weak topology of $H^2(\mu)$. Let us introduce the following numbers

$$\begin{aligned} \alpha_{n,l} &= \int_{1-\delta/l}^1 \frac{1}{2\pi} \int_{-\delta}^{\delta} \frac{r^{2n}}{|1 + \delta/n - re^{it}|^2} dt d\mu(r) \\ &= \frac{2}{\pi} \int_{1-\delta/l}^1 \frac{r^{2n}}{(1 + \delta/n)^2 - r^2} \operatorname{arctg}\left(\frac{1 + \delta/n + r}{1 + \delta/n - r} \operatorname{tg}\left(\frac{\delta}{2}\right)\right) d\mu(r), \end{aligned}$$

and

$$\beta_{n,l} = \frac{2}{\pi} \operatorname{arctg}\left(\frac{1 + \delta/n + 1 - \delta/l}{1 + \delta/n - (1 - \delta/l)} \operatorname{tg}\left(\frac{\delta}{2}\right)\right).$$

Then

$$\begin{aligned} &\left| \left\langle \frac{g_n}{\|g_n\|}, T_f\left(\frac{g_n}{\|g_n\|}\right) \right\rangle \right| \\ &= \left| \frac{1}{\|g_n\|^2} \int_0^1 \int_{\mathbb{T}} f |g_n|^2(rt) d\lambda(t) d\mu(r) \right| \\ &= \frac{1}{2\pi \|g_n\|^2} \left| \int_{1-\delta/l}^1 \int_{-\delta}^{\delta} + \left(\int_0^1 \int_{-\pi}^{\pi} - \int_{1-\delta/l}^1 \int_{-\delta}^{\delta} \right) f |g_n|^2(re^{it}) dt d\mu(r) \right| \\ &\geq \frac{1}{\|g_n\|^2} ((f(1) - \varepsilon)\alpha_{n,l} - \|f\|_{\infty}(\|g_n\|^2 - \alpha_{n,l})) \\ &= (f(1) + \|f\|_{\infty} - \varepsilon) \frac{\alpha_{n,l}}{\|g_n\|^2} - \|f\|_{\infty}. \end{aligned}$$

Applying the fact that functions $r \rightarrow r^{2n}/((1 + \delta/n)^2 - r^2)$ and $r \rightarrow \operatorname{arctg}((1 + \delta/n + r)/(1 + \delta/n - r)\operatorname{tg}(\delta/2))$ are increasing on $[0, 1]$ we get

$$\begin{aligned} \|g_n\|^2 &= \frac{1}{2\pi} \int_{1-\delta/l}^1 \int_{-\delta}^{\delta} + \int_{1-\delta/l}^1 \int_{\delta \leq |t| \leq \pi} + \int_0^{1-\delta/l} \int_{-\pi}^{\pi} |g_n(re^{it})|^2 dt d\mu(r) \\ &\leq \alpha_{n,l} + \int_{1-\delta/l}^1 \frac{r^{2n}(1 - \beta_{n,l})}{(1 + \delta/n)^2 - r^2} d\mu(r) + \frac{(1 - \delta/l)^{2n} \mu([0, 1])}{((1 + \delta/n)^2 - (1 - \delta/l)^2)} \\ &\leq \alpha_{n,l} + (1 - \beta_{n,l}) \frac{\alpha_{n,l}}{\beta_{n,l}} \\ &\quad + \frac{(1 - \delta/l)^{2n} \mu([0, 1])}{((1 + \delta/n)^2 - (1 - \delta/l)^2)} \frac{\alpha_{n,l}}{\int_{1-\delta/2l}^1 r^{2n} \beta_{n,l}/((1 + \delta/n)^2 - r^2) d\mu(r)} \\ &\leq \frac{\alpha_{n,l}}{\beta_{n,l}} \left(1 + \left(\frac{1 - \delta/l}{1 - \delta/2l} \right)^{2n} \frac{\mu([0, 1])}{\mu([1 - \delta/2l, 1])} \right). \end{aligned}$$

Hence for every $l \in \mathbb{N}$

$$\begin{aligned} \lim_n \frac{\alpha_{n,l}}{\|g_n\|^2} &\geq \lim_n \frac{(2/\pi)\operatorname{arctg}(((2 + \delta/n - \delta/l)/(\delta/n + \delta/l))\operatorname{tg}(\delta/2))}{1 + ((1 - \delta/l)/(1 - \delta/2l))^{2n} \mu([0, 1])/\mu([1 - \delta/2l, 1])} \\ &= \frac{2}{\pi} \operatorname{arctg}\left(\frac{2l - \delta}{\delta} \operatorname{tg}\left(\frac{\delta}{2}\right)\right). \end{aligned}$$

Since the sequence $(g_n/\|g_n\|)$ is weakly null in $H^2(\mu)$,

$$\lim_n K\left(\frac{g_n}{\|g_n\|}\right) = 0 \quad \text{for every compact operator } K \in \mathcal{K}(H^2(\mu)).$$

Hence

$$\begin{aligned} \text{dist}(T_f, \mathcal{K}(H^2(\mu))) &\geq \limsup_n \left\| T_f\left(\frac{g_n}{\|g_n\|}\right) \right\| \geq \limsup_n \left| \left\langle \frac{g_n}{\|g_n\|}, T_f\left(\frac{g_n}{\|g_n\|}\right) \right\rangle \right| \\ &\geq (f(1) + \|f\|_\infty - \varepsilon) \frac{2}{\pi} \arctg\left(\frac{2l - \delta}{\delta} \text{tg}\left(\frac{\delta}{2}\right)\right) - \|f\|_\infty. \end{aligned}$$

Since the estimation above holds for every $l \in \mathbb{N}$,

$$\text{dist}(T_f, \mathcal{K}(H^2(\mu))) \geq f(1) - \varepsilon$$

To finish the proof is enough to note that $\varepsilon > 0$ we took arbitrarily. ■

In the sequel we will need the following well known

Fact 4. For every $T \in \mathcal{L}(H^2(\mu))$

$$\begin{aligned} \text{dist}(T, \mathcal{K}(H^2(\mu))) &= \sup\{\limsup_n \|T(h_n)\| : (h_n) \subset B_{H^2(\mu)} \text{ is weakly null}\} \\ &= \lim_n \|T - T \circ P_n\|, \end{aligned}$$

where P_n is the orthogonal projection of $H^2(\mu)$ onto the linear span of e_1, \dots, e_n for some orthonormal Schauder basis (e_n) in $H^2(\mu)$.

Proof. Since

$$\|T - T \circ P_{n+1}\| = \|T \circ (Id - P_n) \circ (Id - P_{n+1})\| \leq \|T - T \circ P_n\|,$$

the sequence $(\|T - T \circ P_n\|)$ is decreasing. The inequality $\text{dist}(T, \mathcal{K}(H^2(\mu))) \leq \lim_n \|T - T \circ P_n\|$ is clear. If $(h_n) \subset B_{H^2(\mu)}$ is a weakly null sequence in $H^2(\mu)$, then

$$\limsup_n \|T(h_n)\| = \limsup_n \|(T - K)(h_n)\| \leq \|T - K\|$$

for every $K \in \mathcal{K}(H^2(\mu))$. Hence

$$\sup_n \{\limsup \|T(h_n)\| : (h_n) \subset B_{H^2(\mu)} \text{ is weakly null}\} \leq \text{dist}(T, \mathcal{K}(H^2(\mu))).$$

Let us select $h_n \in P_n^{-1}(\{0\}) \cap B_{H^2(\mu)} = (Id - P_n)(B_{H^2(\mu)})$ such that

$$\|(T - T \circ P_n)(h_n)\| \geq \|T - T \circ P_n\| - \frac{1}{n}.$$

Then (h_n) is a weakly null sequence in $H^2(\mu)$ and

$$\limsup_n \|T(h_n)\| = \limsup_n \|(T - T \circ P_n)(h_n)\| \geq \lim_n \|T - T \circ P_n\|.$$

■

Straightforward consequence of the theorem and the fact above is the following

Corollary 5. *Let μ be a positive finite Borel measure on $[0, 1]$ such that $1 \in \text{supp}(\mu)$. If $f \in L^\infty(\mu \otimes \lambda)$ is continuous at each point of \mathbb{T} , then*

$$\text{dist}(T_f, \mathcal{K}(H^2(\mu))) = \sup_{t \in \mathbb{T}} |f(t)|.$$

Proof. In view of Theorem 3

$$\text{dist}(T_f, \mathcal{K}(H^2(\mu))) \geq \sup_{t \in \mathbb{T}} |f(t)|.$$

On the other hand, for every $\varepsilon > 0$ there exists $R \in (0, 1)$ such that $\sup\{|f(z)| : z \in \mathbb{D} \setminus R\mathbb{D}\} < \varepsilon + \sup\{|f(t)| : t \in \mathbb{T}\}$. Let $(h_n) \subset B_{H^2(\mu)}$ be a weakly null sequence. Since (h_n) converges uniformly to zero on $R\mathbb{D}$, for every $(g_n) \subset B_{H^2(\mu)}$

$$\begin{aligned} \limsup_n |\langle g_n, f h_n \rangle| &\leq \limsup_n \left(\int_{R\mathbb{D}} |f h_n g_n| d\mu \otimes \lambda \right. \\ &\quad \left. + (\varepsilon + \sup_{t \in \mathbb{T}} |f(t)|) \int_{\mathbb{D} \setminus R\mathbb{D}} |h_n g_n| d\mu \otimes \lambda \right) \\ &\leq \varepsilon + \sup_{t \in \mathbb{T}} |f(t)|. \end{aligned}$$

Hence

$$\text{dist}(T_f, \mathcal{K}(H^2(\mu))) \leq \sup_{t \in \mathbb{T}} |f(t)|. \quad \blacksquare$$

The corollary show that every bounded Borel function f on $\overline{\mathbb{D}}$ with compact support in \mathbb{D} generates the compact Toeplitz operator T_f in each Bergman-Hardy space $H^2(\mu)$. The next result generalizes the following well known fact for Toeplitz operators on the Hardy space H^2 on \mathbb{T} : for every $f \in L^\infty(\lambda)$

$$\text{dist}(T_f, \mathcal{K}(H^2)) = \|f\|_\infty.$$

Recall that for $f \in L^1(\lambda)$ its Poisson integral $\mathbb{P}(f) : \mathbb{D} \rightarrow \mathbb{C}$ is given by

$$\mathbb{P}(f)(z) = \int_{\mathbb{T}} \frac{1 - |z|^2}{|t - z|^2} f(t) d\lambda(t).$$

Every bounded harmonic function F on \mathbb{D} is the Poisson integral of its radial limit function F^* , i.e. $F = \mathbb{P}(F^*)$, and $\sup\{|F(z)| : z \in \mathbb{D}\} = \sup \text{ess}\{|F^*(t)| : t \in \mathbb{T}\}$ (see [2]).

Theorem 6. *Let μ be a positive Borel measure on $[0, 1]$ such that $1 \in \text{supp}(\mu)$. If $f \in L^\infty(\lambda)$, then*

$$\text{dist}(T_{\mathbb{P}(f)}, \mathcal{K}(H^2(\mu))) = \|f\|_\infty.$$

Proof. Since $|\mathbb{P}(f)| \leq \|f\|_\infty$ $\mu \otimes \lambda$ -a.e. on $\overline{\mathbb{D}}$, $\text{dist}(T_{\mathbb{P}(f)}, \mathcal{K}(H^2(\mu))) \leq \|f\|_\infty$. Then we only have to show the inequality in the other direction. Since $\mathbb{P}(f)$ is a bounded and harmonic function on \mathbb{D} , the limit

$$\lim_{r \rightarrow 1} \mathbb{P}(f)(rt) = \mathbb{P}(f)^* \quad \text{exists for } \lambda\text{-a.e. } t \in \mathbb{T}.$$

This limit exists for every Lebesgue point of f (see [2]). Let us take any $\|f\|_\infty > \varepsilon > 0$ and any point t_0 for which the limit above exists and $|\mathbb{P}(f)^*(t_0)| > \|f\|_\infty - \varepsilon$. We can assume that $\mathbb{P}(f)^*(t_0) > 0$. Then there exists $\delta > 0$ such that $|\mathbb{P}(f)(rt_0) - \mathbb{P}(f)^*(t_0)| < \varepsilon$ for every $r \in [(1 - \delta)^3, 1)$. Let

$$g_n(z) = \frac{z^n}{1 + \delta/n - zt_0}.$$

Then

$$\|g_n\|^2 = \int_0^1 \int_{\mathbb{T}} \frac{r^{2n}}{|(1 + \delta/n) - rtt_0|^2} d\lambda(t) d\mu(r) = \int_0^1 \frac{r^{2n}}{(1 + \delta/n)^2 - r^2} d\mu(r).$$

We know from the proof of Theorem 3 that $(g_n/\|g_n\|)$ is a weakly null sequence in $H^2(\mu)$. Let

$$\alpha_n = \int_{1-\delta}^1 \int_{\mathbb{T}} \frac{r^{2n}}{|(1 + \delta/n) - rtt_0|^2} d\lambda(t) d\mu(r) = \int_{1-\delta}^1 \frac{r^{2n}}{(1 + \delta/n)^2 - r^2} d\mu(r)$$

Then

$$\begin{aligned} & \left| \left\langle \frac{g_n}{\|g_n\|}, T_{\mathbb{P}(f)} \left(\frac{g_n}{\|g_n\|} \right) \right\rangle \right| \\ &= \frac{1}{2\pi \|g_n\|^2} \left| \int_{1-\delta}^1 \int_{-\pi}^{\pi} + \int_0^{1-\delta} \int_{-\pi}^{\pi} (\mathbb{P}(f) |g_n|^2)(re^{it}) dt d\mu(r) \right| \\ &\geq \frac{1}{\|g_n\|^2} \left(\left| \int_{1-\delta}^1 \frac{r^{2n}}{(1 + \delta/n)^2 - r^2} \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{(1 + \delta/n)^2 - r^2}{|(1 + \delta/n) - rtt_0|^2} \mathbb{P}(f)(re^{it}) dt d\mu(r) \right| \right. \\ &\quad \left. - \|f\|_\infty (\|g_n\|^2 - \alpha_n) \right) \\ &\geq \frac{1}{\|g_n\|^2} \left(\left| \int_{1-\delta}^1 \frac{r^{2n}}{(1 + \delta/n)^2 - r^2} \mathbb{P}(f) \left(\frac{r^2}{1 + \delta/n} t_0 \right) d\mu(r) \right| - \|f\|_\infty (\|g_n\|^2 - \alpha_n) \right) \\ &\geq \frac{1}{\|g_n\|^2} ((\mathbb{P}(f)^*(t_0) - \varepsilon)\alpha_n - \|f\|_\infty (\|g_n\|^2 - \alpha_n)) \\ &= (\mathbb{P}(f)^*(t_0) + \|f\|_\infty - \varepsilon) \frac{\alpha_n}{\|g_n\|^2} - \|f\|_\infty. \end{aligned}$$

Furthermore

$$\begin{aligned} \|g_n\|^2 &= \frac{1}{2\pi} \int_{1-\delta}^1 \int_{-\pi}^{\pi} + \int_0^{1-\delta} \int_{-\pi}^{\pi} |g_n(re^{it})|^2 dt d\mu(r) \\ &\leq \alpha_n + \frac{(1-\delta)^{2n} \mu([0, 1])}{((1+\delta/n)^2 - (1-\delta)^2)} \frac{\alpha_n}{\int_{1-\delta/2}^1 r^{2n} / ((1+\delta/n)^2 - r^2) d\mu(r)} \\ &\leq \alpha_n \left(1 + \left(\frac{1-\delta}{1-\delta/2} \right)^{2n} \frac{\mu([0, 1])}{\mu([1-\delta/2, 1])} \right) \end{aligned}$$

Hence

$$\lim_n \frac{\alpha_n}{\|g_n\|^2} \geq \lim_n \frac{1}{1 + ((1-\delta)/(1-\delta/2))^{2n} \mu([0, 1]) / \mu([1-\delta/2, 1])} = 1$$

Since $(g_n / \|g_n\|)$ converges weakly to zero in $H^2(\mu)$,

$$\begin{aligned} \text{dist}(T_f, \mathcal{K}(H^2(\mu))) &\geq \limsup_n \left\| T_f \left(\frac{g_n}{\|g_n\|} \right) \right\| \geq \limsup_n \left| \left\langle \frac{g_n}{\|g_n\|}, T_{\mathbb{P}(f)} \left(\frac{g_n}{\|g_n\|} \right) \right\rangle \right| \\ &\geq \mathbb{P}(f)^*(t_0) - \varepsilon \geq \|f\|_{\infty} - 2\varepsilon. \end{aligned}$$

To finish the proof is enough to note that $\varepsilon > 0$ we took arbitrarily. ■

W. Lusky [4] showed that a similar fact holds for the angular extension \tilde{f} of $f \in L^\infty(\lambda)$,

$$\text{dist}(T_{\tilde{f}}, \mathcal{K}(H^2(\mu))) = \|f\|_{\infty}.$$

where $\tilde{f}(rt) = f(t)$ for every $r \in [0, 1)$ and $t \in \mathbb{T}$.

For a bounded holomorphic function f on \mathbb{D} the multiplication operator $M_f : H^2(\mu) \rightarrow H^2(\mu)$ is given by

$$M_f(g) = fg = T_f(g).$$

Multiplication operators on $H^2(\mu)$ form a subclass of Toeplitz operators. Straightforward consequence of the theorem above is the following

Corollary 7. *Let μ be a positive finite Borel measure on $[0, 1]$ such that $1 \in \text{supp}(\mu)$. If f is a bounded holomorphic function on \mathbb{D} , then*

$$\text{dist}(M_f, \mathcal{K}(H^2(\mu))) = \sup_{z \in \mathbb{D}} |f(z)|.$$

Acknowledgments. I wish to thank Professor W. Lusky for many helpful discussions.

References

- [1] L. A. Coburn, *Singular integral operators and Toeplitz operators on odd spheres*, Indiana Univ. Math. J. **23** (1973), 433–439.
- [2] P. L. Duren, *Theory of H^p -Spaces*, Pure Appl. Math. 38, Academic Press, New York 1970.
- [3] Y. Katznelson, *An introduction to Harmonic Analysis*, John Wiley and Sons, New York 1968.
- [4] W. Lusky, *Toeplitz operators on generalized Bergman-Hardy spaces*, Math. Scand., to appear.
- [5] W. Lusky, *On approximation by Toeplitz operators*, Acta Univ. Carolin. Math. Phys. **39** (1998), 137–146.
- [6] G. McDonald and C. Sundberg, *Toeplitz operators on the disc*, Indiana Univ. Math. J. **28** (1979), 595–611.
- [7] A. Michalak, *Translations of functions in vector Hardy classes on the unit disk*, Dissertationes Math. **359** (1996), 1-79.
- [8] K. Stroethoff and D. Zheng, *Toeplitz and Hankel operators on Bergman spaces*, Trans. Amer. Math. Soc. **329** (1992), 773–794.
- [9] K. Zhu, *Operator Theory in Function Spaces*, Marcel Dekker, New York 1990.

Address: Faculty of Mathematics and Computer Sciences, A. Mickiewicz University, Matejki 48/49, 60-769 Poznań, Poland

E-mail: michalak@amu.edu.pl

Received: 15 Jan 2000

